

Non-linear Concentration Effects in Magnetic Particle Imaging

V. C. Behr¹, T. Kampf¹, J.-P. Gehrcke¹, M. A. Rückert^{1,2}, P. Vogel^{1,2}, W. H. Kullmann², and P. M. Jakob^{1,3}

¹Experimental Physics 5, University of Würzburg, Würzburg, Germany, ²Medical Engineering, University of Applied Sciences Würzburg-Schweinfurt, Schweinfurt, Germany, ³Magnetic Resonance Bavaria e.V., Würzburg, Germany

Introduction

The single particle model (SPM) is a well-established ferrofluid magnetization theory in the field of MPI [1,2]. It describes non-interacting magnetic particles, i.e., particle concentrations c close to 0. In a real ferrofluid, particles interact magnetically. The smaller the distance between single magnetic moments, the stronger their interaction. Hence, the magnetization curve $M(H)$ non-linearly depends on c : especially for increasing c magnetization saturates at lower H . For tomographic purposes, the quantity c has to be reconstructed from the MPI signal. Hence, the quality of the theory describing the impact of concentration c on the magnetization curve is essential for image reconstruction. Since the effect described is ignored by the SPM a different model is required to properly fit data.

Simulations

For our simulations, we use a second order modified mean-field theory (MMF2) [3]. MMF2 incorporates magnetic particle coupling and reflects experimental results best of all models tested in [4]. A mono-disperse ferrofluid made of homogeneously distributed magnetite particles at different iron concentrations c_{Fe} is being considered. Using MMF2, the relative magnetization response $M_{rel}(5mT/\mu_0 \sin(\omega_0 t))$ is calculated. Its time derivative is analyzed via discrete Fourier transformation, leading to amplitudes of higher harmonics $A_n(c_{Fe})$. As a consequence of the nonlinear impact of c on $M(H)$, the n -th higher harmonic amplitude A_n in magnetic particle spectroscopy is non-linearly dependent on c , too. As a function of c_{Fe} figure 1 shows the relative magnetization of a sample while figure 2 displays the magnitude of the higher harmonics normalized to the 3rd harmonic are shown. In general, the ratio A_n/A_3 severely increase due to concentration enhancements. Harmonics with higher n are affected more. The SPM is, as expected, not able to exhibit this behavior.




Figure 3: **top:** Phantom $c(x)$. **center:** nonlinear part of the MMF2-simulated signal voltage; acquired during 1 period of drive field trajectory. **bottom:** Reconstructed phantom $c_{reco}(x)$ using different inversion kernels.

substance linearly changes with c_{Fe} . That allowed us to simulate the experimental setup using MMF2 and SPM. The result of the simulations was correlated to the experiment by linear scaling of MMF2 data to experimental data.

The experimental and simulated data is shown in figure 4. The nonlinear $A_n(c)$ dependency could be verified. Furthermore, MMF2 is providing a proper description of the magnetization response of dense ferrofluids.

Conclusions

Higher harmonic amplitudes in MPI are not linearly related to concentration changes, as suggested by the SPM. Therefore, current linear image reconstruction schemes will fail in properly reconstructing spatial particle concentration distributions containing areas with dense ferrofluids. Those will not be uncommon in biological applications of MPI, e.g. the agglomeration of particles in cells – leading to iron concentrations of 0.2 – 5 mol/l in localized areas [6]. Thus, incorporating this non-linear behavior in imaging reconstruction schemes is mandatory. The experimental results suggest that using MMF2 is capable of correctly describing the non-linear effects on the higher harmonics in the MPI signal. The difference between the SPM and MMF2 curves visualizes the inadequacy of the SPM. Therefore it should be used instead of SPM for dense ferrofluids.

Acknowledgements

We acknowledge Prof. Schenk and Reiner Beringer (University of Würzburg, Anorganic Chemistry) for preparation of our ferrofluid samples. This work was supported by grants from the German Federal Ministry of Education and Research (BMBF, FKZ 1745X08) and the German Research Foundation (DFG, SFB 688).

References

- [1] Gleich, Weizenecker, Nature 435 214-7 (2005)
- [2] Rahmer et al., BMC Medical Imaging 9:4 (2009)
- [3] Ivanov, Kuznetsova, Phys. Rev. E 64, 041405 (2001)
- [4] Ivanov et al., Phys. Rev. E 75, 061405 (2007)
- [5] Goodwill et al., IEEE Trans Med Imag Aug;28(8):1231-7 (2009)
- [6] Billotey et al., Magn. Reson. Med. 49:646-654 (2003)




Figure 1: **(top)** Normalized magnetization curves for different concentrations c , calculated using MMF2; in comparison with SPM.

Figure 2: **(bottom)** Normalized signal amplitudes of a spectroscopic MPI experiment in dependence of c . Signal obtained by MMF2 and SPM in comparison.




Figure 4: Experimentally gained amplitudes of the 7th harmonic of the signal of five ferrofluid samples exhibiting different iron concentration c_{Fe} , in comparison with MMF2 and SPM simulations.